



Diabetic Ketoacidosis

Overview

Make the diagnosis
Establish venous access
Fluids
Insulin
Monitoring
Underlying cause

NB There is no difference in the management of DKA in a new or established patient

Definition

Blood glucose > 11 mmol/L
pH < 7.3
Bicarbonate < 15 mmol/L
AND
more than 5% dehydrated
and/or vomiting
and/or drowsy
and/or clinically acidotic

Features of DKA

Polyuria, polydipsia, incontinence
Thirst, polydipsia
Abdominal pain
Vomiting
Kussmaul breathing
Acetone on breath
Dehydration, hypotension, collapse
Disturbed consciousness
Coma

Features of shock

Poor capillary return
Impaired consciousness
BP may be normal or low
Thready, rapid pulse

NB Urine output may remain fair because of osmotic diuresis

Features of Cerebral Oedema

Headache
Irritability
Slowing of pulse
Rising blood pressure
Reducing conscious level

NB These features usually present a few hours after commencing treatment but can occur very early

Biochemical Assessment and Monitoring

Blood

All new diabetes patients

Glucose

U&Es including Bicarbonate

Thyroid function tests

FBC

HbA1c

10ml plain tube for possible autoantibodies
(HISS Order Set /NEWDIA)

for DKA

Gases (venous usually sufficient)

Blood culture

Osmolality

Urine

All patients

Glucose, Ketones (test all urine and record results)

for DKA

Hourly volume, culture



Other monitoring etc in DKA

Measure **current weight** if at all possible

BP

ECG monitor

Nil by mouth

NG Tube if unconscious or evidence of gastric distention

Neuro-obs

Repeat biochemistry frequently (~2hrly) until patient stable

Fluids for DKA - use with DKA IV Fluid Calculation Sheet



1. Treat shock with 10ml/kg N Saline. May repeat once before discussing with a consultant.
2. Calculate fluid deficit - see A, B, C and D on calculation sheet
3. Determine maintenance fluid volume per kg from age:

0-2 years	160ml per kg
3-5 years	140ml per kg
6-9 years	120ml per kg
10-14 years	100ml per kg
>14 years	60ml per kg

NB Maintenance calculation is for 48 hours

Weigh the child and compare with previous known weights and centile charts - access the electronic medical record for existing patients

4. Calculate hourly fluid rate (for 48h)

$$\text{Hourly Rate (ml)} = \frac{[48\text{h MAINTENANCE (C x E)} + \text{DEFICIT (D x 1000)}] - \text{Volume used to treat shock}}{48}$$

5. Double check all calculations and have someone else do so independently - if you are in any doubt **ASK**

DKA Treatment - Which Fluids?



General Points:

Treat shock (resuscitate) with Normal Saline

Sodium Bicarbonate is very rarely indicated and may be harmful - only consider in Intensive Care Unit after discussion with consultant - dose would be 0.5mmol/kg over 30 minutes

Potassium can usually be added to bags immediately after resuscitation (assuming urine output) - 20 mmol/500ml. Monitor T waves and adjust KCl according to electrolyte results. Insulin makes K fall.

Phosphate - it is normal for patients in DKA to have a low phosphate level as this is a plasma buffer. There is no evidence to support adding additional phosphate.

Initially use 0.9% saline.

Generally, once the blood glucose has fallen to 14 mmol/l add glucose to the fluid.

If this occurs **within** the first 6 hours, the child may still be sodium depleted. Discuss this with consultant, who may wish to continue with Normal saline and added dextrose.

If this occurs **after** the first 6 hours and the child's plasma sodium level is stable, change the fluid type to 0.45% saline/5% dextrose.

After 1st 12h, and assuming that the patient is improving, if they have already changed to 0.5N Saline + 5% Dextrose, there is no need to change back to Normal Saline if glucose > 14mmol/L - **ADJUST the INSULIN**



DKA Treatment - Insulin

Insulin is essential to switch off ketogenesis

Make up a solution of 1 unit per ml. of human soluble insulin (e.g. Actrapid) by adding 50 units (0.5 ml) insulin to 49.5 ml 0.9% saline in a syringe pump. Attach this using a Y-connector to the IV fluids already running.

Do not add insulin directly to the fluid bags.

Run at 0.1 units/kg/hour (0.1ml/kg/hour).

If the rate of blood glucose fall exceeds 5 mmol/l per hour, or falls to around 14 mmol/l, add dextrose (5-10% equivalent) to the IV fluids running (see “fluids” above). The insulin dose needs to be maintained at 0.1 units/kg/hour to switch off ketogenesis.


Do not stop the insulin infusion while dextrose is being infused, as insulin is required to switch off ketone production. If the blood glucose falls below 4 mmol/l, give a bolus of 2 ml/kg of 10% dextrose and increase the dextrose concentration of the infusion.

10% dextrose with 0.45% saline can be made up by adding 7.5ml NaCl 30% to 500ml 10% Dextrose. (remember to consider if KCl required)

Once the pH is above 7.3, the blood glucose is down to 14-17 mmol/l, and a dextrose-containing fluid has been started, consider reducing the insulin infusion rate, but to no less than 0.05 units/kg/hour.

Blood glucose rises out of control, or the pH level is not improving after 4-6 hours consult senior medical staff, re-evaluate (possible sepsis, insulin errors or other condition), and consider starting the whole protocol again.

DKA Treatment - Cerebral Oedema



Symptoms and Signs:

Headache, slowing of heart rate, rise in BP
Change in neurological status - restlessness, irritability, drowsiness, incontinence.
Specific neurological signs e.g. cranial nerve palsies
Abnormal posturing

Highest risk 12-18h after beginning rehydration

Inform senior staff immediately

Treat in ICU

Document carefully

Management:

Exclude hypoglycaemia as a possible cause of any behaviour change

Give Mannitol 1 g/kg stat (= 5 ml/kg Mannitol 20% over 20 minutes) or hypertonic saline (5-10 mls/kg over 30 mins). This needs to be given as soon as possible if warning signs occur.

Restrict IV fluids to 2/3 maintenance and replace deficit over 72 rather than 48 hours

Discuss with PICU consultant (if assisted ventilation is required maintain pCO₂ above 3.5 kPa)

Once the child is stable, exclude other diagnoses by CT scan - other intracerebral events may occur (thrombosis, haemorrhage or infarction) and present similarly

A repeated dose of Mannitol should be given after 2 hours if no response

DKA Intravenous Fluids Calculations

For all calculations for paediatric DKA please go to ggc-youngdiabetes.org and look under DKA